



VACCINATION POLICY

PKC DOGGY DAY CARE AND CATTERY

Vaccinations need to be current and must be given at least one week prior to coming into kennels. If vaccinations have not been given regularly and they are overdue, there may be a longer stand down period required so please contact the kennels if you are unsure.

Please scan through the vaccination book to allow us to sight current vaccination status of your animals.

Cats:

We require vaccinations to be annual to come into our cattery.

There are three types of core vaccines available (all cats must have one of these every year)

1. Felocell 3 / 4
2. Tricat Trio
3. Nobivac Ducat

Protocol for cat vaccines (same for all types):

INITIAL COURSE (kittens or previously unvaccinated cats, 2 or 3 doses, Kittens between the ages of 6-16 weeks, or

adults). Boosters 3 – 4 weeks apart, no more than 6 weeks.

FIRST ANNUAL: An additional shot is required either at 12 months of age or 12 months after the last shot, this

completes the initial course

BOOSTERS: After the initial course, annual vaccinations are required

Dogs:

We require all dogs to have up to date:

(A). Core Vaccinations, and (B). Canine Cough protection

A. 5 in 1 (core vaccine) - There are two main brands of CORE VACCINES (all dogs must have) available to cover Parvovirus, Distemper and Adenovirus:

- | | |
|------------------|-----------|
| 1). VANGUARD 5 / | 2). DHP / |
| VANGUARD PLUS 5 | DHPPI |

Protocol for VANGUARD 5: (Usually Adult Dogs)

INITIAL COURSE: Two vaccinations, Initially, Boosters 3 – 4 weeks apart

BOOSTERS: Adults: Annual required

Protocol for VANGUARD PLUS 5: (Usually Puppies)

INITIAL COURSE: Three vaccinations, 3 - 4 weeks apart from 6 weeks of age, (16 week finish)

FIRST ANNUAL: 12 – 16 months: One Shot

BOOSTERS: Adults: One Shot up

Protocol for DHPPI and DHP:

INITIAL COURSE: Puppies: 2 – 3 vaccinations – last dose can be given at 10 weeks, or over 3 – 4 weeks apart

FIRST ANNUAL: 12 – 16 months: One Shot

BOOSTERS: Adults: One Shot every 3 years

B. Canine Cough Vaccine - There are three types of CANINE COUGH vaccines – all dogs must have one of these annually

Protocol for Injectable CC:

INITIAL COURSE: (Puppies or new dogs): Two vaccines 3 – 4 weeks apart, from 6 weeks

12 – 16 months: One Shot

BOOSTERS: Annual boosters required

Protocol for Intranasal CC:

INITIAL COURSE: Puppies: One vaccination 8 weeks or older

BOOSTER: 12 – 16 months: One Shot

BOOSTER: Adults: Annual boosters required

Protocol for Oral KC:

One vaccination from 8 weeks, then annually

Please note: if your dog has had Intranasal vaccines every year but the most recent CC vaccine was Canigen KC or Canvac BB, if they haven't had a booster of this injection 3-4 weeks later, they are incorrectly vaccinated and you need to contact your Vet to find out what to do.

NO EXCEPTIONS